

Full Length Research Paper

Sensory assessment of sorghum brew adjunct and barley brew lager beer

Olu Malomo*, Ogunmoyela O. A. B., Oluwajoba S. O., Adigun M. O. and Daniel 'Toyosi

College of Food Science, Bells University of Technology, Ota, Ogun State, Nigeria.

Accepted 19 October, 2011

The type and strain of yeast used in fermentation has a great influence on the taste and character of beer produced. Apart from brewing, sorghum has been used extensively in food industries. The essence of this study is to investigate the behavior of yeast in a sorghum/barley brew and also to investigate the sensory acceptability of such a combination. The study showed that the assessors were unable to detect if there were differences below a 40/60 blend, sorghum and barley mix respectively, beyond which all the organoleptic parameters presented became objectionable.

Key words: Organoleptic, fermentation, assessors, brewing, sorghum, barley.

INTRODUCTION

Alcoholic beverages which occur throughout the world in many forms and taste results from the action of micro-organisms or enzymes on a wide range of agricultural products such as grapes, grains, and soybeans (Smith, 1996; Rose, 1989). Such biological action was associated with biochemical changes that gave rise to significant organoleptic improvements to the final products (Brill, 1981). These products which are more nutritious and more easily digestible are toxicologically and microbiologically safer (Smith, 1996). The process of fermenting raw materials is of wide diversity, using technology from the most primitive to the most advanced achieving an outstanding range of sensory and textural quality in the final product (Rose, 1989). Alcoholic beverages and potable spirit industries represent one of the most economically stable sectors in the present day commerce. Materials for alcoholic beverages normally comprise either sugary material (fruit juice, plant sap) or starchy materials (grains or roots) which need to be hydrolyzed to simple sugar before fermentation. When these substrates are incubated with suitable micro-organism and allowed to ferment, the end product is a liquid containing from a few percentages to about 16% alcohol with a slightly acidic pH.

The type of alcoholic beverage produced in any particular region or country almost entirely reflects the types of crop grown. Thus the cooler region of Europe, Scandinavia, Poland and Russia produces beer and lagers from barley (Palmer, 1992). In tropical Africa, alcoholic beverages especially beer have been brewed for generations with locally available cereals like Sorghum, Rice, Maize and Millet (Okafor, 1987).

Barley malt is also known to be rich in protein and enzymes. It has a high β -glucan and pentosan level when compared to local grains. The enzyme potential of barley malt is sufficient to catabolize additional starch. Consequently, throughout the world, part of the malt, usually 15-20% is replaced by unmalted cereal. This unmalted cereal called adjunct act to remove excess protein which could cause haziness if it finds its way into beer (Kunze, 1996).

Barley malt, whose diastatic power is higher than other, cereals, has a high level of maltose producing β -amylase which is the key enzyme in breaking down malt starch (Palmer, 1996). Barley contains on dry weight basis as shown in (Table 1). Due to economic reasons, cheap raw materials are used as adjunct as partial replacement for malt in wort production (Canales, 1979). In some African countries and in some parts of U.S.A, sorghum has been the grain of choice as adjunct in lager beer production (Okolo and Lewis, 1995).

Sorghum is the fifth largest cereal produced in the world. It is drought resistant and indigenous to Africa. It

*Corresponding author. E-mail: oludaremalomo1951@yahoo.com.

Table 1. Gravity value, pH, color, temperature and attenuation limit of substrate after brewing (24 h)

Malt /Adjunct Ratio.	pH	Color (EBC)	Gravity (°P)	Temperature (°C)	Attenuation
100% Barley malt	4.98	5.25	11.0	10.5	2.0
90% barley: 10% sorghum	4.87	5.25	11.0	11.2	2.0
80% barley: 20% sorghum	4.48	5.75	11.2	10.8	2.0
70% barley: 30% sorghum	4.86	6.50	11.3	11.5	2.0
60% barley: 40% sorghum	5.00	7.00	11.5	13.2	2.0

is a staple food rating third behind rice and wheat (Okolo and Lewis, 1995). It belongs to the grass family, essentially adapted to the semi-arid conditions. Where it is traditionally used as food, beverages and building materials (Gomez, 1992). Sorghum known as *Sorghum vulgare* is an old crop with yet unknown genetic variability (Palmer, 1992). It is a dry land summer cereal cultivated in areas of rainfall of 400 and 750 mm and a temperature of 38-40°C (House, 1993).

During germination, sorghum produced an enzyme, which breaks down the storage materials it contains. Since planting is during the raining season, contamination by microorganisms especially fungi are to expect. It is necessary to treat harvested crop with fungicide to prevent spoilage (Gomez, 1992). Sorghum has a wide range of colors and sizes. The best known are the white (*farafara*), yellow (*kaura*) brown and the red types (Gomez, 1992). Although the red types are malted for traditional beer production, the white type is preferred for modern beer production (Palmer, 1992).

The type and strain of yeast used in fermentation has a great influence on the taste and character of beer produced (Kunze, 1996). Apart from brewing, sorghum has been used extensively in food industries. Vitamin enriched sorghum based product are enjoying acceptance at institutional levels as mid shift drinks and high energy breakfast food (Aluko, 1989). Sorghum is also used in industrial production of cellulose, paper, starch, sugar and chemicals (Subramanian and Jambunathan, 1989).

Other uses of sorghum include livestock feed, production (Al-Hazzan et al., 1989). Sorghum as adjunct in beer production already has a widespread acceptance. However, the availability of the grain poses a great problem to the brewing industry. Sorghum is already a staple diet, so to make a substantial replacement for barley malt, huge quantities of adjunct will be required (Nweke and Ibe, 1989).

In brewing the basic principles remain the same over the centuries. This includes malting, mashing, boiling and fermentation (Kunze, 1996). During malting, the grains are allowed to steep in water for 48-72 h at 10-15°C. The grains are allowed to germinate. However, as soon as the enzymes are produced and before the young seedling can make an appreciable in road into the nutrient reserve of the grain, development of the seedling is halted by

drying at a temperature which will not completely inactivate the enzymes.

These enzymes are reactivated during mashing to hydrolyze starch and protein and release nutrients for the nourishment of the yeast during fermentation (Okafor, 1987). After malting the grains are milled and mixed with water. The mash after several processes of brewing is filtered and boiled before fermentation. Wort is usually inoculated with the fermenting organism, yeast. The sugar in the wort is fermented by yeast to alcohol. For this purpose yeast specie such as *Saccharomyces carlsbergensis* are used. Selected strains are systematically isolated and grown (Pollock, 1981).

Yeast has a high vitamin and enzyme content. They are particularly rich in vitamins B₁ and B₆ (Smith, 1996). Yeast makes use of carbohydrates in two ways: by respiration and fermentation. During fermentation, the nitrogen content of the wort, formed as a result of protein hydrolysis should at least be 23 mg of free amino nitrogen to allow for proper yeast nutrition (Kunze, 1996). Yeast cells are usually ovoid or round in shape.

They are non-motile cells measuring between 5 x 10⁶ (5 nm) and 10 x 10⁶ (10 nm) in diameter. They reproduced by budding and grow rapidly on a defined medium. They also have a well-developed genetic cycle (Pollock, 1981). Yeast cells are able to ferment the trisaccharide raffinose, made up of glucose, fructose, and galactose. When using an adjunct as a replacement for malt during brewing, it must be remembered that protein is deposited in the adjunct in a more stable form. The protein is degraded only to a slight extent, because of this; adjunct mashes contain less molecular weight nitrogenous materials (alpha amino nitrogen) than malt mashes. Adjunct beer therefore always contains more polyphenols. It follows then that the higher the adjunct the less the nitrogenous compounds (Kunze, 1996).

MATERIALS AND METHODS

Raw materials used for brewing were collected from the warehouse of the International Breweries PLC Ilesa. Barley malt (*Hordeum distichon*) or (*Hordeum vulgare*) was the chief raw material used. Others include, raw sorghum (*Sorghum vulgare*) was used as adjunct, brewing water, yeast (*Saccharomyces carlsbergensis*), hop, and other additives like Vitamin C (ascorbic acid), sulphuric acid, sodium hydroxide and calcium chloride. These additives are used to adjust the pH of wort, when and where necessary.

Sample collection

Pilsener barley malt (*Hordeum disichon*) with moisture content of 4.55%, kernel weight of 33.4 g was used as the substrate in brewing. The barley was collected from the warehouse of International Breweries PLC, Ilesa. The yellow type raw sorghum (*Sorghum vulgare*) with a moisture content of 6.15% and kernel weight of 35.5 gm was used as adjunct. This was purchased from L.O. Omole and Sons store, Ilesa.

Brewing process

A laboratory manual Glasbaserei milling machine was used to reduce the particle sizes of the grains and also to expose the starch in the grains in readiness for enzymatic action during brewing. In 100% barley malt brew, 600gm of malted barley was weighed into a flask. In the case of other formulations, the following proportions were used; 90% barley: 10% sorghum = 540 gm barley: 60gm sorghum. 80% barley: 20% sorghum = 480 gm barley: 120gm sorghum. 70% barley: 30% sorghum = 420 gm barley: 180gm sorghum. 60% barley: 40% sorghum = 360 gm barley: 240 gm sorghum.

Two 25 l capacity Carlsberg's flask were used for fermentation. An electronic julabo water bath was used to monitor the temperature of wort during brewing.

Procedure

Six hundred grams of malted barley was weighed and mashed with water inside a conical flask. This mash was placed inside a water bath and the temperature was raised to between 50 and 55°C for about 20 min, to allow for optimum activity of the enzymes present in the malt. The temperature was later raised to 72°C for 45 min. to terminate enzymatic activities. The mash was then allowed to rest for 45 min to allow the mash to undergo complete saccharification (Okafor, 1987).

Saccharification, which is the complete hydrolysis of starch to simple sugar is shown by a brown or colorless reaction with 2% iodine solutions (Brauhaase, 2000). After saccharification, the mash was filtered. The aqueous solution collected that is, wort was boiled for 1 h. After boiling, the wort was allowed to cool and the following parameters were determined; pH, Color, gravity and temperature.

The cooled wort was transferred into the Carlsbergen's vessel and inoculated with the fermenting organism. The pitching wort was then kept in the refrigerating incubator. With malt/ adjunct brew the gelatinized sorghum adjunct was added to the malt mash at 55°C. The enzyme in the barley malt also acted on the adjunct mash to hydrolyze the gelatinized starch in the adjunct.

Fermenting organism

A genetically modified strain of *Saccharomyces carlsbergensis* no. 01185, Strain 370 from Horlunann, Hamburg, was used as the fermenting organism. The yeast used was obtained by the propagation of pure yeast culture in the Laboratory.

Yeast propagation

About 5 ml of sterile wort was poured into a test-tube, a loopful of yeast from PDA agar slant was used to inoculate the sterile wort and shaken vigorously. The test tube was allowed to stand for 24h. The inoculated wort was transferred into a 100 ml flask containing sterile wort and this was also allowed to stand for another 24 h. This was also transferred into another flask containing 250 ml of

sterile wort and subsequently to 500 and 1000 ml sterile wort and each was allowed to stand for 24 h. At the end 1000 ml of fermenting wort was obtained and this was used as the inoculant for the fermentation process.

Fermentation

Wort collected after brewing was transferred into Carlsberg's vessels. About 150 ml of the propagated inoculant with a yeast count of about 22 million cells / ml was introduced into the fermenting vessels using an injection syringe. The flask was shaken vigorously and kept in the refrigerating incubator with an inserted thermometer to monitor the temperature. Fermentation was allowed to proceed for nine days and the pH, Temperature, Color, Gravity, Yeast count, Attenuation, Yeast viability and Yeast consistency were monitored and determined daily.

pH determination

The pH was determined using a Jenway 3015 pH meter. Ten milliliter of fermenting wort was taken and the electrode of the pH meter was inserted into the wort sample. The reading on the screen of the pH meter was observed and recorded daily. The pH value of wort after brewing i.e. first day of fermentation is found in Table 1.

Color determination

The color of wort before and after fermentation was taken recorded. The wort sample was put in a color comparator (colorimeter). The color of the wort at 320 nm wavelength was matched with the color in the color comparator and the values recorded. The color of wort before fermentation was recorded in Table 1.

Yeast viability and consistency

About 5 mL of fermenting wort was placed in a test tube, two drops of methylene blue was added to the sample, the sample was vigorously shaken and one or two drops of the stained sample was placed on the haemocytometer, mounted on the microscope and observed. The dead cells which absorbed the stain and retained the blue stain of the methylene blue was expressed as the percentage of the living cells which absorbed the blue color of the methylene blue and digested it to become colorless (Peppler, 1978). Yeast viability is calculated as follows:

$$\frac{\text{Total number of cells} - \text{Dead cells}}{\text{Total number of cells}} \times 100$$

Yeast consistency was carried out by placing about 20 mL of fermenting wort in a cuvette. The cuvette was placed in the chamber of a Heraeus Christ Labofuge centrifuge. The sample in the cuvette was spinned at 4000 rpm. After spinning, the supernatant was discarded and the compressed yeast was weighed and recorded. This experiment was carried out in duplicates and the result was calculated as follows (Rose, 1977):

$$\frac{\text{Tube 1} = e - a}{c - a} \times 100 \%$$

$$\frac{\text{Tube 2} = f - b}{d - b} \times 100 \%$$

Table 2. Parameters monitored during fermentation: temperature, pH, attenuation limit, (apparent) and deformed yeast.

Days of fermentation	Temperature (°C)					Ph					Attenuation limit (fermented extract)					Deformed yeast				
	100	90:10	80:20	70:30	60:40	100	90:10	80::20	70::30	60:40	100	90:10	80:20	70:30	60:40	100	90:10	80:20	70:30	60:40
Day 1	13.5	12.0	10.2	12.6	10.8	4.98	4.96	4.43	4.86	5.00	2.0	2.0	2.4	2.0	2.0	-	-	-	-	-
Day 2	12.8	12.2	12.0	10.8	10.8	4.80	4.90	4.33	4.49	4.63	7.82	14.54	28.6	18.6	16.03	-	-	-	-	-
Day 3	10.8	12.6	11.9	10.8	11.4	4.51	4.64	4.21	4.35	4.20	12.73	21.81	32.14	34.5	33.0	-	-	-	-	-
Day 4	10.6	12.4	11.8	10.7	12.0	4.21	4.52	4.18	4.32	3.73	18.26	41.86	69.6	50.4	59.8	-	-	-	-	-
Day 5	10.4	11.4	11.6	10.6	10.2	4.10	4.40	4.02	4.44	3.70	34.78	45.4	79.5	60.17	70.4	-	-	-	-	-
Day 6	10.4	10.6	10.8	10.2	10.2	4.08	4.08	4.02	4.26	3.75	44.34	47.2	80.4	72.6	82.6	-	-	-	-	-
Day 7	9.0	10.0	8.6	8.6	10.0	4.06	4.06	4.00	4.30	3.76	63.47	78.2	81.25	77.9	84.3	-	-	-	-	-
Day 8	8.5	8.2	7.5	7.5	8.2	4.02	4.04	3.88	4.32	3.77	82.6	83.6	81.25	85.0	87.0	-	-	-	-	+
Day 9	6.2	6.5	5.0	6.8	5.0	4.0	4.0	3.80	4.15	3.72	84.34	89.0	82.14	89.3	89.56	-	-	-	-	+

Table 3. Original gravity, final gravity, specific gravity, real extract, refractive index, and alcohol of fermented extracts.

Proportions	Original gravity	Final gravity	Specific gravity	Real extract	Refractive index	Alcohol
100% Barley malt	11.5	2.0	1°°/602	3.84	36.5	3.95
90% Barley : 10% sorghum	11.0	1.8	1°°/702	3.95	36.45	3.61
80% Barley : 20 % sorghum	11.2	2.1	1°°/820	4.20	37.38	3.90
70% Barley : 30% sorghum	11.3	1.7	1°°/663	3.80	36.50	3.78
60% Barley : 40 % sorghum	11.5	1.5	1°°/585	3.75	36.35	3.90

Where: Weight of tube 1 = a; Weight of tube 2 = b; Weight of tube 2 + fermenting wort = d; Weight of tube 1 + fermenting wort = c; Weight of tube 1 + compressed yeast = e and Weight of tube 2 + compressed yeast = f.

Attenuation determination

The attenuation was calculated by expressing the final gravity (Fg) as a percentage of the original gravity (Og), that is;

$$\frac{Og - Fg}{Og} \times 100\%$$

After fermentation, the green beer was filtered to remove the yeast cells. The clear beer was then checked for final gravity, pH, color and alcohol. The alcohol content was determined using the Variag Hans Carl alcohol chart. On this chart, the Original gravity (Og) and final gravity (Fg) of the beer was used to determine the specific gravity (Sg), real extract (Re), refractive index (RI) and alcohol content of the beer for each formulations (Kunze,1996) (Table 3).

RESULTS AND DISCUSSION

pH

In the different formulations, it was observed that

pH was between 4.48 in 80% barley: 20% sorghum to 5.00 in 60% barley: 40% sorghum. These pH values still conform to the desired pH range of wort, which is between 4.0 and 5.4 (Kunze, 1996), Table 1 shows the pH values of the wort.

Colour (EBC –European Brewery Convention)

The color of the wort increased with increase in adjunct concentration from 5.25 in 100% barley malt wort to 7.00 in 60% barley: 40% sorghum

adjunct (Table 1). This could be attributed to the color of raw sorghum used due to the presence and oxidation of polyphenols and melanoidin in the sorghum (Brauhaase, 2000).

Gravity

The gravity of the wort increased with increase in the adjunct proportion (Table 1). This implies that sorghum adjuncts contains more fermentable sugar than barley malt due to the high starch content of sorghum. Moreover very little of the soluble protein is contributed to the wort (about 5% of their total protein) (Brauhaase, 2000).

Temperature

The temperature of fermentation was maintained at between 10 to 13.2°C. This is to allow for initial vigorous fermentation. A low initiating temperature may lead to too sluggish fermentation due to cold shock that could be experienced by the yeast cells on pitching while a higher temperature would lead to too vigorous fermentation thereby producing unwanted by product (Hough et al., 1977).

Attenuation

The attenuation value of all the formulations was recorded in Table 1. All values are within the specified limit. The essence of attenuation is to know at what value the primary fermentation should be terminated so that enough extract will be available for secondary fermentation in the storage tanks. The attenuation values of all the formulation ranged between 2.0 –2.4 as recorded in Table 2.

Brewing

During brewing process, it was observed that the brew with 40% sorghum adjunct took a longer period of time to saccharify (30 min) as compared to the brews with lower adjunct proportion (8-10 min). It was also observed that the filtration rate of the high adjunct proportion brew was slower than the other formulations. This could as a result of the endosperm hardness of the sorghum grains (Malomo, 1993).

Fermentation

During fermentation, the green beer in all the formulations was covered with white layers of foam. As fermentation progressed, the bubbles became fluffy and

form brown caps. The appearance of bubbles showed that fermentation was progressing favorably. Towards the end of fermentation, the high crest of foam formed slowly collapsed and the foam appeared browner and the bubbles became less pronounced. This showed that fermentation process had slowed down, less carbon dioxide was been produced and fermentation was slowly going to an end (Kunze, 1996).

Gravity

The amount of fermentable sugars reduced as fermentation progressed. This is because the yeast cells utilized these sugars which were converted to ethyl alcohol and carbon dioxide. The gravity of wort dropped almost 24 h of primary fermentation. This is because about 3% of the extract were fermented every 24 h (Brauhaase, 2000). The gravity fall corresponded with an increase in the yeast population. This is because yeast cells multiply when the nutrients was available in the medium. An increase in yeast population indicated that more yeast cells utilized the sugar in the wort more readily, thereby bringing down the gravity of the wort. The peak of the yeast population was recorded on the 4th day of fermentation in all cases of formulations except the 60%:40% malt adjunct proportion. This inconsistency could be as a result of early depletion of the nutrients in the substrate. It is known that the higher the adjunct, proportion, the lower the availability of free amino nitrogen in the substrate. This is because the protein in the adjunct is deposited in the crude (unhydrolysed) form and so the amino acid in the form of α - amino nitrogen is not readily available for yeast nutrition (Kunze, 1996).

In all the other formulations, after the 4th day of fermentation, yeast population dropped gradually. The metabolic activity of the cells reduced drastically due to the depletion of nutrients in the medium, eventually resulting in the settling down of the yeast cells called yeast flocculation. However, further drop in gravity was observed until the gravity of the wort was between 1.2 – 2.0°P on the 9th day of fermentation (that is, the final gravity (FG)). The original gravity of the wort (gravity on the first day of fermentation) ranged between 11.0 – 11.5°P.

Temperature

The temperature of the fermenting medium ranged between 10.2 to 13.2°C. The temperature of the fermenting wort was maintained at between 10.8 to 12.8°C. This was to allow the yeast cells to act at their optimum temperature. As yeast cells act well at this temperature range (Pollock, 1981). It should be noted that the temperature in the fermenting vessel was controlled by a cooling device (refrigerating incubator),

because metabolic activity of yeast cells would lead to temperature increase inside the fermenting liquor which would lead to an increase in fermentation rates and therefore reduces fermentation period. The cumulative effect of this action is the production of unwanted by-products such as more diacetyl. The foam and colloidal stability of the beer may worsen. From the 6th day of fermentation, a drop in temperature was observed. This was because the metabolic activity of the yeast had greatly reduced. The low temperature observed also assisted in yeast flocculation (Kunze, 1996).

pH

The pH of the pitching wort ranged between 4.43 –5.00. Ideally, the pH of any pitching wort is between 4.5 –5.6 (Goldammer, 2002). As fermentation progressed, the pH decreased until the final day (day 9). The pH values ranged between 3.72 in 60: 40 malt/adjunct to 4.15 in 70:30 malt adjunct ratio. The reduction in pH values of the fermenting liquor could be as a result of the production of organic acids, uptake of ammonium ions, and the use of primary phosphate ions by the yeast (Kunze, 1996). A drastic fall in pH observed in the 60:40 malt/adjunct ratio is not desirable in the beer because it imparts acidic taste to the beer (Hough, 1977). This low pH value also affected the growth and performance of yeast cells in the fermenting wort. Generally speaking, the pH desired by any brewer is between 4.2 and 4.4 (Brauhaase, 2000).

The fluctuation in the pH in the fermenting wort on days 5 and 7 of the 70:30 malt/adjunct ratio and in 60:40 malt/adjunct ratio on days 6, 7 and 8 could be as a result of yeast autolysis. This results from the process of self digestion of the yeast cells when there is an early depletion of nutrients in the wort (Kunze, 1996).

Yeast count

Yeast count increased progressively in all generations as observed from day 2 to day 4 of fermentation. It followed a normal growth curve whereby growth increased exponentially after introducing a microbe into a nutrient until limiting growth factors bring it down (Pollock, 1981). In this fermentation, the yeast in the 60:40 malt/adjunct proportions was not consistent. The yeast dropped after day 2 of fermentation. This could be due to the high percentage of adjunct and because of the low value of the free amino nitrogen in the wort, essential nutrient likes, amino acids, lysine, and nitrogen were not available for proper yeast nutrition and nourishment.

Alcohol content

The desired alcohol content of beer is generally $3.92 \pm$

0.08% (Pollock, 1981). All formulations however produced beer of desired alcohol level. Considering the high extract fermented in 60:40 malt/adjunct ratios (Table 3), it was expected that the alcohol content should be higher than other formulations (Table 3). This could be related to the fact that most of the available nutrients in the medium had been used up for yeast nutrition and nourishment rather than for alcohol production. Table 3 shows the original gravity, final gravity, specific gravity, real extract, refractive index and alcohol content of all the formulations.

Apparent attenuation (fermented real extract)

Apparent attenuation is the total amount of fermented extract in the wort. From the experiment, it was observed that the highest quantity of fermentable extract was recorded in the 60:40 malt /adjunct formulation (56°P) while the least observed was recorded in 80:20 malt /adjunct (82.14°P). The high extract percentage used up by yeast in the 0:40 malt /adjunct formulation showed that most of the extract had been utilized by yeast during primary fermentation. This would undoubtedly have a negative effect on the secondary fermentation, which is the maturity stage of the beer (Anon, 1972). Table 2 shows the values of fermented extract for each formulation.

Conclusion

All proportions of malt/ adjunct ratio produced beer of desirable physico chemical parameters however 60:40 malt adjunct ratio produced beer with low pH which could have deleterious effect on the beer and the yeast cells used in fermentation. The emergence of deformed yeast after fermentation showed that the 60:40 formulations did not contain enough nutrients for yeast nutrition and metabolism.

Since the aim of every brewing outfit is to be able to compete favorably in the highly competitive Nigerian beer market as well as to produce beer of optimum quality, it is advisable to stick to beer produced with malt /adjunct proportion of up to 70:30 malt /adjunct proportion whose yeast behavior can be scientifically predicted to avoid deleterious side effects like fusel oil production, acid, sulfide and other volatile compounds which could affect the quality of the beer. Also good beer yeast is expected to be used up to 8 – 10 generations (Pollock, 1981). The emergence of deformed yeast in the 60:40 malt / adjunct ratio in the first generation showed that the yeast used might not be able to withstand the stress of long usage.

However the use of external enzymes like amylase, protease and filtrase could assist in bringing out optimum activity when using sorghum adjunct in a higher proportion as in 60:40 malt adjunct proportion. The use of higher percentage of adjunct should therefore be

explored putting in consideration the use of external enzymes which although would give desired end product but will be more finance intensive.

REFERENCES

- Al-Hazzan, Umunna NN, Alawa JP (1989). Symposium, at the International crop research institute for the semi-arid tropics (INCRISAT).
- Aluko AO (1989). Cereal Science today. Rev Biotechnol., 12:259.
- Anon NA (1972). Alcoholic Beverages. Academic Press. New York. pp. 139.
- Brauhaase (2000). Standard operating manual. General parameters in brewing. pp. 3-49.
- Brill LW (1981). Cereal science today. Food Sci. Technol. Report, 3 .7ed. pp. 259.
- Canales AM (1979). Unmalted grains in brewing science. Vol. 1 Academic press, London. pp. 146-23.
- Gomez MI (1992). Screening of sorghum for malting and use. Agriculture of sorghum symposium. (SADCC/INCRISAT). Bulawayo. Zimbabwe
- Hough JS, Briggs DE, Stevens R (1977). Malting and Brewing Science. Chapman and Hall London. Pp. 8.
- Hough JS (1977). Development of brewing Analysis. A Historical review. The Institute of Brewing. London. pp. 12.
- House LR (1993). Agriculture of sorghum. Symposium for the International crop research institute for the semi- arid tropics. (SADCC/ICRISAT).
- Kunze W (1996). Technology malting and brewing. VLB Germany. pp. 120-250, 253-259.
- Malomo O (1993). Sorghum in lager beer production in Nigeria. Regional symposium on current progress in the processing and utilization of sorghum and other related cereals. Ouagadougou, Burkina Faso.
- Nweke FI, Ibe DG (1989). Nigeria Agriculture and Brewing; A new symbiosis. Symposium Brew. Ind., pp. 13- 16.
- Okafor N (1987). Industrial Microbiology. 1st edition University of Ife Press. pp.174 - 182.
- Okolo BN, Lewis IE (1995). Enhancement of amyolytic potentials of sorghum malt. J. Inst. Brew., 102:267-274.
- Palmer GH (1992). Sorghum, The world class cereal, Symposium on science and technology, Belgium.
- Pollock JRA (1981). Brewing science. Vol. 2 Academic press New York and London. pp. 491.
- Rose AH (1989). The microbial production of food and drinks. The advent of genetic engineering and food Microbiology. A handbook of Microbiology. pp. 45-54.
- Smith EY (1996). Introduction to Biotechnology. Academic press, New York, 1st Edition. pp. 185-186.
- Subramanian V, Jambunathan R (1989). Industrial utilization of sorghum. Symposium on International crop research institute for the semi arid tropics. (ICRISAT).